This presentation should have included all images analyzed in the paper plus some images (from CG/Chat-Cre-ZsGreen) mice for another project Due to size limit, it is printed to pdf.

Rationale:


Previous report in cancer research indicated that VEGFR1 activation let to increased thermal and mechanical sensitivity via TRPV-1 and TRPA-1 mediated pathways, respectively.

Western blot analysis showed that VEGFR1, but not VEGFR2, was significantly upregulated in the lysates of ipsilateral L3–L4 lumbar DRGs isolated from tumor-bearing mice as compared with sham-treated animals.


Question: Does the expression level of VEGFRs, TRPV1 and TRPA1 change in DRG sensory neuronal somata in response to bladder VEGF_A instillation?


Experimental design: Littermates receive bladder instillation of either VEGF_A or saline. Immunostaining for VEGFR1, R2, TRPV-1, and TRPA1 in L5/L6-S2 DRG.

The 488 channel was reserved for ZsGreen signal, 594 channel was used for anti-VEGFR1, anti-VEGFR2, or anti-TRPV-1. DAPI was used to reveal the cell nuclei.




Antibody against VEGFR1 exhibits broad staining in DRG neurons and potentially satellite glia cells (1), with strong perinuclear staining (2). The VEGFR1 signal is also detected in the cytosol of small diameter neurons (3).






20190328 RGs/PBSVR1 RG female_R0165_40x_1 Saline Instillation


*Threshold is set based on non-primary images from the same experiments (day - should have been animals)




VEGF R1 Staining day 1




VEGF R1 Staining day 2







VEGF R1 Staining day 4 CGs



VEGFR1 Signal in both sexes



*Normalized VEGFR1 Signal in both sexes




Total count
 36
 M
 F


 Saline
 17
 9
 8

 VEGF165
 19
 9
 10


^{*} Normalized to the average intensity of the salineinstilled tissue from the same staining day.

Two schools of thoughts:








VEGFR-2 is robustly expressed by endothelial cells throughout the extensive DRG capillary network, but not found in sensory neurons or other nonendothelial cells of the DRG.




D Freshly dissected










VEGF R2 Staining day 2








Normalized VEGFR2 Signal in both sexes




Normalized VEGFR2 Signal; combined sexes





Total count	31	M	F
Saline	14	7	7
VEGF165	17	9	8


TRPV1 Staining Saline Instillation - 1



RG-F-R2378 No primary









TRPV1 Staining VEGF_A Instillation - 2

Normalized TRPV-1 Signal in both sexes




Normalized TRPV-1 Signal; combined sexes




Total count	21	M	F
Saline	11	6	5
VEGF165	10	5	5

Antibody against TRPA1 exhibits signal in satellite glia cells and some DRG neurons.




Antibody against TRPA1 exhibits signal in satellite glia cells and some DRG neurons.



20190328 RGs/PBSVR1 RG female_R0165_40x_1 Saline Instillation

Normalized TRPA-1 Signal in both sexes



Normalized TRPA-1 Signal; combined sexes

